

CLIMATE SUMMARY FOR: SKEENA REGION

PART OF A SERIES ON THE RESOURCE REGIONS OF BRITISH COLUMBIA

ABOUT THIS SERIES

There is a strong scientific consensus that the Earth's climate is changing, primarily due to greenhouse gas emissions. This series of climate summaries, for the eight resource regions of British Columbia, is meant to help inform readers about past climate and future projected changes. It is intended that the series will be updated with new information as research progresses.

GENERAL OVERVIEW

British Columbia's climate exhibits large variations over short distances, due to complex topography.

Long-term historical trends show warming, more rapid for night-time low temperatures than daytime highs and more rapid in winter than summer. Precipitation trends are less certain due to data limitations and also exhibit increases, except in the winter season when large variability results in trends that depend highly on the period considered.

Further warming and precipitation changes are projected throughout the 21st century. The magnitude of the projected warming is relatively large compared to historical variability. Some possible consequences of these projected changes on resource operations are considered.

ABOUT THIS REGION

The Skeena Region, with a population of just under 90,000, is in the northwestern corner of British Columbia, stretching south from the Yukon to Haida Gwaii and encompassing the inland area east of the Alaska Panhandle (Figure 1). The region contains parts of the Skeena and Liard drainage basins. Owing to the complex topography, which includes parts of the Coast, Cassiar, St. Elias and Skeena Mountains as well as the Yukon and Stikine Plateaus, the area's climate varies considerably over short distances. Also, two major Pacific climate patterns—El Niño and the Pacific Decadal Oscillation—exert their influence over the region's year-to-year variability.




Figure 1: Winter precipitation for the region. The region is bounded in grey and the red box shows its location in BC.

The region has both coastal and interior western hemlock and western redcedar forests. Mountain hemlock, alpine communities and glaciers occur at higher elevations on the coast. The interior part of the region has white and black spruce and lodgepole pine dominated forests with Engelmann spruce subalpine fir at higher elevations. Spruce, willow and birch forests and alpine communities occupy the drier, northern part of the region. The economy is largely based around fishing, forestry, energy, transport and tourism.



Precipitation is historically greatest in the autumn and winter seasons, and least in the spring and summer. Precipitation varies considerably throughout the region. Areas with the least winter precipitation historically (less than 200 mm) include the areas in the rainshadow of the Coast Mountains. By contrast, the western edge of the region includes several locations with winter precipitation over 500 mm and some with well over 1000 mm.

HISTORICAL TRENDS

The historical annual trend (based on the CANGRID dataset¹) indicates that close to 2 °C of warming has already occurred during the 20th century. Summer and winter trends are plotted in Figures 2 and 3, while trends for all seasons are provided in Tables 1 and 2. These trends




Figure 2: Historic summer and winter temperature time series, 1901-2009.

Table 1: Temperature Trends (°C per decade) for the Skeena Region

| Period | Trend* 1901-2009 | Statistical Uncertainty in Trend ³ 1901-2009 | Trend* 1951-2009 | Statistical Uncertainty in Trend ³ 1951-2009 |
|--------------|---------------------|---|---------------------|---|
| Spring (MAM) | 0.21 | 0.11 to 0.30 | 0.27 | 0.00 to 0.52 |
| Summer (JJA) | 0.16 | 0.11 to 0.20 | 0.17 | 0.05 to 0.30 |
| Autumn (SON) | 0.08 | 0.01 to 0.15 | 0.09 | -0.11 to 0.28 |
| Winter (DJF) | 0.22 | 0.06 to 0.40 | 0.41 | -0.01 to 0.82 |
| Annual | 0.16 | 0.10 to 0.23 | 0.21 | 0.05 to 0.38 |

*The reported trend is the trend that best describes the change over time in the observations. **Bold** indicates a trend that is statistically significant at the 5% significance level. Multiply the trend by 5 or 10 to get the total amount of change over a 50 or 100-year period, respectively.

are regional averages. In regions with complex topography, trends could vary considerably with elevation.

Warming has occurred in all seasons, more rapidly since the middle of the 20th century. In all seasons 1901-2009 trends are large relative to historical variability, as indicated by statistical significance.

The historical mean seasonal precipitation for the region is greatest in the autumn (about 200 mm), with larger precipitation amounts in mountainous terrain along the eastern border of the region. Precipitation also varies considerably from year to year within the region, as shown in Figure 3.

Precipitation in the region has been increasing over both time periods during all seasons, with the exception of the 1951-2009 winter




Figure 3: Historic summer and winter precipitation time series, 1901-2009.

Table 2: Precipitation Trends (mm/season per decade) for the Skeena Region

| Period | Trend* 1901-2009 | Statistical Uncertainty in Trend ³ 1901-2009 | Trend* 1951-2009 | Statistical Uncertainty in Trend ³ 1951-2009 |
|--------------|---------------------|---|---------------------|---|
| Spring (MAM) | 1 | 0 to 3 | 2 | -3 to 7 |
| Summer (JJA) | 3 | 1 to 5 | 5 | -1 to 10 |
| Autumn (SON) | 5 | 1 to 8 | 0 | -8 to 8 |
| Winter (DJF) | 4 | 0 to 7 | -5 | -14 to 4 |

*The reported trend is the trend that best describes the change over time in the observations. **Bold** indicates a trend that is statistically significant at the 5% significance level. Multiply the trend by 5 or 10 to get the total amount of change over a 50 or 100-year period, respectively.

precipitation trend, which is negative and the autumn trend which is zero. Low confidence in precipitation observations in the early part of the century implies a need for caution in interpreting the difference between short- and long-term winter precipitation trends. Large year-to-year and decade-to-decade variability in winter precipitation and the choice of time period used for fitting trends also affect this result.

FUTURE CLIMATE PROJECTIONS

Climate models project⁴ warming throughout the 21st century for all seasons that is large compared to historical variability (Figure 4). The black bar shows historical interannual variability as represented by \pm one standard deviation of temperature around the 1961-1990 average (vertical line). The projected change in the average is shown for three future periods.

Projected warming is quite uniform across the seasons, with annual warming of 1.8 °C (1.1 °C to 2.5 °C) by the 2050s and 2.6 °C (1.5 °C to 4.3 °C) by the 2080s.

Projected precipitation changes are relatively modest compared to historical variability, as shown in Figure 4. By the 2080s the median projection indicates an increase of about 15 %, relative to the 1961-1990 baseline, in all seasons but summer.

Note that in Table 3 and Figure 4, the projections from two different emissions scenarios (A2 and B1) are combined to give a range of anticipated future change. In the early and middle of the 21st century, the emissions scenario has little influence on the amount of projected change. The ensemble projected annual warming is 2.6 °C (1.5 °C to 4.3°C) by the 2080s. The projections following the higher (A2) emissions scenario represent the warmer portion of the projected range of change (and vice versa for lower emissions, B1).




Figure 4: Cumulative seasonal precipitation and mean seasonal temperature projections for three future periods, the 2020s (2011-2040), 2050s (2041-2070) and 2080s (2071-2100). These are 30-year regional averages. The width of the bands indicate the range of projections. The thin, upper black line and the lower band indicate the average and the variability, respectively, over the 1961-1990 reference period.

The summer mean temperature for the Skeena region during the 20th century was about 10 °C. The warmest 10% of summers were almost 2 °C warmer than this average, about 12 °C averaged across the entire region. Under the median summer warming of 2.5 °C, about two-thirds of summers in the 2080s would be warmer than the warmest 10% of summers in the past even if no change in the distribution of temperature extremes occurs.

SUMMARY OF PROJECTED CHANGE

Table 3 is from Plan2Adapt.ca, a PCIC product that provides projections for the 21st century, as well as interactive maps and information on impacts.

By the 2050s, there are substantial projected decreases in spring snowfall, and a decrease in heating degree days. Along with these changes, an increase in frost-free days and growing degree days is indicated.

POTENTIAL IMPACTS

Changes to the overall climate of the region can result in a variety of associated impacts. This section makes use of Plan2Adapt's impacts tab, which displays impacts that could potentially be associated with the amount of temperature and precipitation change projected for the region.

Adapting forests to the region's climate changes will likely require increasing species diversity and using assisted migration. A change in agricultural productivity could result from a longer growing season, seasonally waterlogged soil and decreased water availability. New crops and varieties may become viable.

Warming and an accompanying reduction in snowpack could result in a shorter winter logging season.

Warming will decrease snowpack throughout much of the region. A seasonal increase in hot and dry conditions would decrease the supply of water. An increase in rain on snow, and high-intensity precipitation can increase freeze-thaw processes, decrease slope stability and increase the risk of landslides.

Increases to high-intensity precipitation and seasonal moisture variability could have effects on a variety of habitats. Both river flooding frequency and runoff may increase; stream bank erosion and strain on flood protection infrastructure may increase; new design guidelines for flood control levels and infrastructure may be required. Seasonal water quality may be reduced. Water sources for irrigation may become salinated. Stormwater design standards may no longer be adequate.

1. CANGRID is a historical gridded data set with a spatial resolution of 50 km based on station observations, composed by Environment Canada (Zhang et al., 2000: Temperature and precipitation trends in Canada during the 20th century. *Atmosphere Ocean*, **38**, 395-429.).
2. These values are from the PRISM data set, the details of which are given in: Daly, C., et al., 2008. Physiographically-sensitive mapping of temperature and precipitation across the conterminous United States. *International Journal of Climatology*, **28**, 2031-2064.
3. The statistical uncertainty indicates the range of trend estimates that are plausibly consistent with the year-to-year variation in seasonal means. This range is calculated as a statistical "95 % confidence interval."
4. The projected change given is the median from an ensemble of 30 global climate model projections from the Coupled Model Intercomparison Project 3 (CMIP3). The range, in brackets, is the 10th to 90th percentile of projected changes. Details about the ensemble, known as PCIC30, are given in: Murdock, T. Q. and D. L. Spittlehouse, 2011: Selecting and Using Climate Change Scenarios for British Columbia. Pacific Climate Impacts Consortium, University of Victoria, Victoria, British Columbia.

Table 3: Summary of Climate Projections for the Skeena Region

| Climate Variable | Season | Projected Change from 1961-1990 Baseline | |
|---|--------|--|--------------------------|
| | | Ensemble Median | Range (10th-90th %ile) |
| Mean Temperature, 2050s (°C) | Annual | +1.8 °C | +1.1 °C to +2.5 °C |
| Precipitation, 2050s (%) | Annual | +7% | +3% to +13% |
| | Summer | +2% | -5% to +11% |
| | Winter | +9% | -1% to +16% |
| Snowfall*, 2050s (%) | Winter | -6% | -12% to +7% |
| | Spring | -56% | -68% to -10% |
| Growing Degree Days*, 2050s (degree days) | Annual | +226 degree days | +142 to +353 degree days |
| Heating Degree Days*, 2050s (degree days) | Annual | -645 degree days | -918 to -418 degree days |
| Frost-Free Days*, 2050s (days) | Annual | +22 days | +12 to +34 days |

The table above shows projected changes in average (mean) temperature, precipitation and several derived climate variables from the baseline historical period (1961-1990) for the 2050s. The ensemble median is a mid-point value, chosen from a PCIC standard set of Global Climate Model (GCM) projections (Murdock and Spittlehouse 2011). The range values represent the low and high results within the set. Further information, including projections for the 2020s and 2080s see www.Plan2Adapt.ca.

* Derived from temperature and precipitation.